

IMM - DTU

02405 Probability
2003-10-17
BFN/bfn

We let X_i denote the lifetime of the i 'th component,

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t)$$

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t)$$

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t}) (1 - e^{-\lambda_2 t})$$

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t}) (1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t)$$

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t}) (1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t}) (1 - e^{-\lambda_2 t})$$

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317.

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t)$$

IMM - DTU

02405 Probability

2003-10-17

BFN/bfn

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

Question c) From the system design we deduce $S = \max(\min(X_1, X_2), \min(X_3, X_4))$ such that

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

Question c) From the system design we deduce $S = \max(\min(X_1, X_2), \min(X_3, X_4))$ such that

$$P(S \leq t)$$

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

Question c) From the system design we deduce $S = \max(\min(X_1, X_2), \min(X_3, X_4))$ such that

$$P(S \leq t) = (1 - e^{-(\lambda_1 + \lambda_2)t})(1 - e^{-(\lambda_3 + \lambda_4)t})$$

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

Question c) From the system design we deduce $S = \max(\min(X_1, X_2), \min(X_3, X_4))$ such that

$$P(S \leq t) = (1 - e^{-(\lambda_1 + \lambda_2)t})(1 - e^{-(\lambda_3 + \lambda_4)t})$$

Question d) Here $S = \min(\max(X_1, X_2), X_3)$ such that

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

Question c) From the system design we deduce $S = \max(\min(X_1, X_2), \min(X_3, X_4))$ such that

$$P(S \leq t) = (1 - e^{-(\lambda_1 + \lambda_2)t})(1 - e^{-(\lambda_3 + \lambda_4)t})$$

Question d) Here $S = \min(\max(X_1, X_2), X_3)$ such that

$$P(S \leq t)$$

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

Question c) From the system design we deduce $S = \max(\min(X_1, X_2), \min(X_3, X_4))$ such that

$$P(S \leq t) = (1 - e^{-(\lambda_1 + \lambda_2)t})(1 - e^{-(\lambda_3 + \lambda_4)t})$$

Question d) Here $S = \min(\max(X_1, X_2), X_3)$ such that

$$P(S \leq t) = 1 - (1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}))e^{-\lambda_3 t}$$

We let X_i denote the lifetime of the i 'th component, and S denote the lifetime of the system.

Question a) From page 316 and example 4 page 317/318 we have the maximum of two exponential random variables $S = \max(X_1, X_2)$.

$$P(S \leq t) = P(\max(X_1, X_2) \leq t) = (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t})$$

Thus

$$P(S > t) = 1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}) = e^{-\lambda_1 t} + e^{-\lambda_2 t} - e^{-(\lambda_1 + \lambda_2)t}$$

Question b) In this case we have $S = \min(X_1, X_2)$ and we apply the result for the minimum of random variables page 317. The special case of two exponentials is treated in example 3 page 317

$$P(S \leq t) = 1 - e^{-(\lambda_1 + \lambda_2)t}$$

Question c) From the system design we deduce $S = \max(\min(X_1, X_2), \min(X_3, X_4))$ such that

$$P(S \leq t) = (1 - e^{-(\lambda_1 + \lambda_2)t})(1 - e^{-(\lambda_3 + \lambda_4)t})$$

Question d) Here $S = \min(\max(X_1, X_2), X_3)$ such that

$$P(S \leq t) = 1 - (1 - (1 - e^{-\lambda_1 t})(1 - e^{-\lambda_2 t}))e^{-\lambda_3 t} = 1 - e^{-(\lambda_1 + \lambda_3)t} - e^{-(\lambda_2 + \lambda_3)t} + e^{-(\lambda_1 + \lambda_2 + \lambda_3)t}$$