

IMM - DTU

02405 Probability
2003-11-12
BFN/bfn

Question a) The probability $P(X_i \leq x) = x$

IMM - DTU

02405 Probability

2003-11-12

BFN/bfn

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed.

IMM - DTU

02405 Probability

2003-11-12

BFN/bfn

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$.

IMM - DTU

02405 Probability

2003-11-12

BFN/bfn

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

IMM - DTU

02405 Probability

2003-11-12

BFN/bfn

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x)$$

IMM - DTU

02405 Probability
2003-11-12
BFN/bfn

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k)$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $\text{bin}(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $\text{bin}(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution.

IMM - DTU

02405 Probability
2003-11-12
BFN/bfn

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x)$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $\text{bin}(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $\text{beta}(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $\text{beta}(r, s)$ density is

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $beta(r, s)$ density is

$$f(x)$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $beta(r, s)$ density is

$$f(x) = \frac{1}{B(r, s)} x^{r-1} (1-x)^{s-1}$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $beta(r, s)$ density is

$$f(x) = \frac{1}{B(r, s)} x^{r-1} (1-x)^{s-1} = \frac{1}{B(r, s)} x^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-x)^i$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $beta(r, s)$ density is

$$f(x) = \frac{1}{B(r, s)} x^{r-1} (1-x)^{s-1} = \frac{1}{B(r, s)} x^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-x)^i$$

Now

$$P(X_{(k)} \leq x)$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $beta(r, s)$ density is

$$f(x) = \frac{1}{B(r, s)} x^{r-1} (1-x)^{s-1} = \frac{1}{B(r, s)} x^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-x)^i$$

Now

$$P(X_{(k)} \leq x) = \int_0^x f(x) dx$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $bin(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $beta(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $beta(r, s)$ density is

$$f(x) = \frac{1}{B(r, s)} x^{r-1} (1-x)^{s-1} = \frac{1}{B(r, s)} x^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-x)^i$$

Now

$$P(X_{(k)} \leq x) = \int_0^x f(x) dx = \int_0^x \frac{1}{B(r, s)} u^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-u)^i du$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $\text{bin}(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $\text{beta}(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $\text{beta}(r, s)$ density is

$$f(x) = \frac{1}{B(r, s)} x^{r-1} (1-x)^{s-1} = \frac{1}{B(r, s)} x^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-x)^i$$

Now

$$\begin{aligned} P(X_{(k)} \leq x) &= \int_0^x f(x) dx = \int_0^x \frac{1}{B(r, s)} u^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-u)^i du \\ &= \frac{1}{B(r, s)} \sum_{i=0}^{s-1} \int_0^x \binom{s-1}{i} (-u)^{r+i-1} du \end{aligned}$$

Question a) The probability $P(X_i \leq x) = x$ since X_i is uniformly distributed. Since the X_i are independent, the number N_x of X_i 's less than or equal to x follows a binomial distribution $\text{bin}(n, x)$. The event $\{X_{(k)} \leq x\}$ corresponds to $\{N_x \geq k\}$. We get

$$P(X_{(k)} \leq x) = P(N_x \geq k) = \sum_{i=k}^n \binom{n}{i} x^i (1-x)^{n-i}$$

Question b) From the boxed result at the bottom of page 327 we have that $(X_{(k)})$ has $\text{beta}(k, n - k + 1)$ distribution. Substituting $r = k$ and $s = n - k + 1$ we get

$$P(X_{(k)} \leq x) = \sum_{i=r}^{r+s-1} \binom{r+s-1}{i} x^i (1-x)^{s+r-i-1}$$

which is the stated result.

Question c) The $\text{beta}(r, s)$ density is

$$f(x) = \frac{1}{B(r, s)} x^{r-1} (1-x)^{s-1} = \frac{1}{B(r, s)} x^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-x)^i$$

Now

$$\begin{aligned} P(X_{(k)} \leq x) &= \int_0^x f(x) dx = \int_0^x \frac{1}{B(r, s)} u^{r-1} \sum_{i=0}^{s-1} \binom{s-1}{i} (-u)^i du \\ &= \frac{1}{B(r, s)} \sum_{i=0}^{s-1} \int_0^x \binom{s-1}{i} (-u)^{r+i-1} du = \frac{x^r}{B(r, s)} \sum_{i=0}^{s-1} \binom{s-1}{i} \frac{(-x)^i}{r+i} \end{aligned}$$

as was to be proved.