

IMM - DTU

02405 Probability  
2006-11-28  
BFN/bfn

Question a) We define  $U$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test.

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454.

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300)$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000)$$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(V - 1300\right)$$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90}\right)$$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \middle| U\right)$$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid U - 1200\right)$$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100}\right)$$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page

IMM - DTU

02405 Probability

2006-11-28

BFN/bfn

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2)$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5)$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457.

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates.

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0)$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0)$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0)$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360}$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0)$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5}$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

we get

Question a) We define  $U$  to be a student's score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

we get

$$P(1300 + 90 \cdot Y - (1200 + 100 \cdot X) > 50)$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

we get

$$P(1300 + 90 \cdot Y - (1200 + 100 \cdot X) > 50)$$

$$= P(1300 + 90 \cdot (\rho X + \sqrt{1 - \rho^2}Z) - (1200 + 100 \cdot X) > 50)$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

we get

$$\begin{aligned} &P(1300 + 90 \cdot Y - (1200 + 100 \cdot X) > 50) \\ &= P(1300 + 90 \cdot (\rho X + \sqrt{1 - \rho^2}Z) - (1200 + 100 \cdot X) > 50) \\ &= P(72Z - 46X \geq -50) \end{aligned}$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

we get

$$\begin{aligned} &P(1300 + 90 \cdot Y - (1200 + 100 \cdot X) > 50) \\ &= P(1300 + 90 \cdot (\rho X + \sqrt{1 - \rho^2}Z) - (1200 + 100 \cdot X) > 50) \\ &= P(72Z - 46X \geq -50) = 1 - \Phi\left(-\frac{50}{\sqrt{46^2 + 72^2}}\right) = \end{aligned}$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

we get

$$\begin{aligned} &P(1300 + 90 \cdot Y - (1200 + 100 \cdot X) > 50) \\ &= P(1300 + 90 \cdot (\rho X + \sqrt{1 - \rho^2}Z) - (1200 + 100 \cdot X) > 50) \\ &= P(72Z - 46X \geq -50) = 1 - \Phi\left(-\frac{50}{\sqrt{46^2 + 72^2}}\right) = \Phi(0.585) = \end{aligned}$$

Question a) We define  $U$  to be a students score on the PSAT test and  $V$  to be the score of the same student on the PSAT test. The pair  $(U, V)$  follows a general bivariate normal distribution as given in the box at the bottom of page 454. The probability in question is  $P(V > 1300|U = 1000)$  which we rewrite

$$P(V > 1300|U = 1000) = P\left(\frac{V - 1300}{90} \mid \frac{U - 1200}{100} = -2\right)$$

Now using the definition on page 454 together with the definition of the standard bivariate normal distribution page 451 we get

$$P(0.6 \cdot X + \sqrt{1 - 0.6^2}Z > 0|X = -2) = P\left(Z > \frac{1.2}{0.8}\right) = 1 - \Phi(1.5) = 0.0668$$

Question b) The solution to this question is closely related to the method of Example 2 page 457. First we realize that we can consider standard normal variates. Using the notation of the previous question we formulate the problem as

$$P(Y > 0|X < 0) = P(0.6 \cdot X + 0.8 \cdot Z > 0|X < 0) = \frac{P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0)}{P(X < 0)}$$

Now using the rotational symmetry we see that

$$P(0.6 \cdot X + 0.8 \cdot Z > 0, X < 0) = \frac{90 - \tan^{-1}\left(\frac{3}{4}\right)}{360} = 0.14758. \text{ Finally}$$

$$P(Y > 0|X < 0) = \frac{0.14758}{0.5} = 0.2952.$$

Question c) We formulate the question using the notation of question a) as

$$P(V - U > 50)$$

we get

$$\begin{aligned} &P(1300 + 90 \cdot Y - (1200 + 100 \cdot X) > 50) \\ &= P(1300 + 90 \cdot (\rho X + \sqrt{1 - \rho^2}Z) - (1200 + 100 \cdot X) > 50) \\ &= P(72Z - 46X \geq -50) = 1 - \Phi\left(-\frac{50}{\sqrt{46^2 + 72^2}}\right) = \Phi(0.585) = 0.72 \end{aligned}$$