

Geostatistics and Analysis of Spatial Data

Allan A. Nielsen

Abstract— This note deals with geostatistical measures for spatial correlation, namely the auto-covariance function and the semi-variogram, as well as deterministic and geostatistical methods for spatial interpolation, namely inverse distance weighting, radial basis functions (RBF) and kriging. Some semi-variogram models are mentioned, specifically the spherical, the exponential and the Gaussian models. Equations for RBF interpolation as well as simple and ordinary kriging (OK) are deduced. Other types of kriging are mentioned, and references to international literature, Internet addresses and state-of-the-art software in the field are given. A very simple example to illustrate the computations for OK and a more realistic example with height data from an area near Slagelse, Denmark, are given. A series of attractive characteristics of kriging are mentioned, and a simple sampling strategic consideration is given based on the dependence of the kriging variance of distance and direction to the nearest observations.

I. INTRODUCTION

Often we need to be able to integrate point attribute information with vector and raster data which we may already have stored in a Geographical Information System (GIS). This can be done by linking the point information to a geographical coordinate in the data base. If we have lots of point data, a tempting alternative will be to generate an interpolated map so that from our point data we calculate raster data which can be analysed along with other sources of raster data.

This note deals with geostatistical methods for description of spatial correlation between point measurements as well as deterministic and geostatistical methods for spatial interpolation.

The basic idea in geostatistics consists of considering observed values of geochemical, geophysical or other natural variables as realisations of a stochastic process in the 2-D plane or in 3-D space. For each position \mathbf{r} in a domain \mathcal{D} which is a part of Euclidian space, a measureable quantity $z(\mathbf{r})$ termed a *regionalised variable* exists. $z(\mathbf{r})$ is considered as a realisation of a *stochastic variable* $Z(\mathbf{r})$. The set of stochastic variables $\{Z(\mathbf{r}) \mid \mathbf{r} \in \mathcal{D}\}$ is termed a *stochastic function*. $Z(\mathbf{r})$ has mean value or expectation value $E\{Z(\mathbf{r})\} = \mu(\mathbf{r})$ and auto-covariance function $\text{Cov}\{Z(\mathbf{r}), Z(\mathbf{r} + \mathbf{h})\} = C(\mathbf{r}, \mathbf{h})$, where \mathbf{h} is termed the displacement vector. If $\mu(\mathbf{r})$ is constant over \mathcal{D} , i.e., $\mu(\mathbf{r}) = \mu$, Z is said to be first order stationary. If also $C(\mathbf{r}, \mathbf{h})$ is constant over \mathcal{D} , i.e., $C(\mathbf{r}, \mathbf{h}) = C(\mathbf{h})$, Z is said to be second order stationary.

This statistical view is inspired by work carried out by Georges Matheron in 1962-1963. It is described in for example [1], [2], [3] gives a good practical and data analytically oriented introduction to geostatistics. [4] is a chapter in a collection of articles which describe many different techniques and their application within the geosciences. [5] deals with geostatistics and other relevant subjects in the context of analysis of spatial data. Geostatistical expositions in a GIS context can be found in [6], [7]. [8] deals with multivariate geostatistics, i.e., studies of the joint spatial co-variation of more variables. The International Association for Mathematical Geology (IAMG) publishes i.a. the periodical *Mathematical Geosciences* (formerly *Mathematical Geology*) where many results on geostatistical research are published. *State-of-the-art open source software* may be found in *GSLIB*, [9], and *Variowin*, [10]. Other easily obtainable softwares are *Geo-EAS* and *Geostatistical Toolbox*. All these pack-

ages can be found at <http://www-sst.unil.ch/research/variowin/> (or via a search engine). Also commercial geostatistical software exists.

This note which is inspired by [11] (see also [12]), in Section II deals with spatial correlation, specifically the auto-covariance function, the semi-variogram and some semi-variogram models are described. Section III deals with spatial interpolation including the deterministic methods inverse distance weighting and radial basis function interpolation along with a family of statistically based methods termed kriging. Here simple and ordinary kriging are dealt with in some detail. Section IV gives final remarks.

II. SPATIAL CORRELATION

This section mentions methods for description of similarity between measurements of natural variables in the 2-D plane or in 3-D space. Specifically the auto-covariance function and the semi-variogram are introduced. Also a relation between the two is given.

A. The Semi-Variogram

Consider two scalar quantities $z(\mathbf{r})$ and $z(\mathbf{r} + \mathbf{h})$ measured at two points in the plane or in space \mathbf{r} og $\mathbf{r} + \mathbf{h}$ separated by the displacement vector \mathbf{h} . We consider z as a realisation of a stochastic variable Z . The variability may be described b.m.o. the *auto-covariance function* (assuming or enforcing first order stationarity, i.e., the mean value is position independent)

$$C(\mathbf{r}, \mathbf{h}) = E\{[Z(\mathbf{r}) - \mu][Z(\mathbf{r} + \mathbf{h}) - \mu]\}.$$

The *variogram*, 2γ , is defined as

$$2\gamma(\mathbf{r}, \mathbf{h}) = E\{[Z(\mathbf{r}) - Z(\mathbf{r} + \mathbf{h})]^2\},$$

which is a measure for the expected squared difference between stochastic variables as a function of position and the displacement vector. In general the variogram will depend on the displacement vector \mathbf{h} as well as on the position vector \mathbf{r} . The *intrinsic hypothesis* of geostatistics says that the *semi-variogram*, γ , is independent of the position vector and that it depends only on the displacement vector, i.e.,

$$\gamma(\mathbf{r}, \mathbf{h}) = \gamma(\mathbf{h}).$$

If $Z(\mathbf{r})$ is second order stationary (i.e., its auto-covariance function is position independent), the intrinsic hypothesis is valid whereas the opposite is not necessarily true.

If we assume or enforce second order stationarity the following relation between the auto-covariance function and the semi-variogram is valid

$$\gamma(\mathbf{h}) = C(\mathbf{0}) - C(\mathbf{h}). \quad (1)$$

Note, that $C(\mathbf{0}) = \sigma^2$, the variance of the stochastic variable.

Given a set of point measurements the semi-variogram may be calculated b.m.o. the following estimator, which calculates (half) the mean value of the squared differences between all pairs of measurements $z(\mathbf{r}_k)$ and $z(\mathbf{r}_k + \mathbf{h})$ separated by the displacement vector \mathbf{h}

$$\hat{\gamma}(\mathbf{h}) = \frac{1}{2N(\mathbf{h})} \sum_{k=1}^{N(\mathbf{h})} [z(\mathbf{r}_k) - z(\mathbf{r}_k + \mathbf{h})]^2.$$

$N(\mathbf{h})$ is the number of point pairs separated by \mathbf{h} . $\hat{\gamma}$ is termed the *experimental semi-variogram*. Often we calculate mean values of $\hat{\gamma}$ over intervals $\mathbf{h} \pm \Delta\mathbf{h}$ for both length (magnitude) and direction (argument) of \mathbf{h} . Mean values for the magnitude of \mathbf{h} ($h \pm \Delta h$) are calculated in order to get a sufficiently high $N(\mathbf{h})$ to obtain a low estimation variance for the semi-variogram value. Mean values over intervals of the argument of \mathbf{h} are calculated to check for possible anisotropy. Anisotropy refers to the characteristic that the auto-covariance function and the semi-variogram do not behave similarly for all directions of the displacement vector between observations. This possible anisotropy may also be checked by calculating 2-D semi-variograms also known as variogram maps, [3], [13], [11], [10], [9].

B. Semi-Variogram Models

In order to be able to define its characteristics we parameterise the semi-variogram b.m.o. different semi-variogram models. An often used model, γ^* , is the *spherical model* (here we assume or impose isotropy, i.e., the semi-variogram depends only on distance and not on direction between observations, and we denote by h the magnitude of \mathbf{h})

$$\gamma^*(h) = \begin{cases} 0 & h = 0 \\ C_0 + C_1 \left[\frac{3}{2} \frac{h}{R} - \frac{1}{2} \frac{h^3}{R^3} \right] & 0 < h < R \\ C_0 + C_1 & h \geq R, \end{cases}$$

where C_0 is the so-called *nugget effect* and R is termed the *range of influence* or just the *range*; $C_0/(C_0+C_1)$ is the relative *nugget effect* and C_0+C_1 is termed the *sill* ($= \sigma^2$). The parameters C_0 and C_1 are not to be confused with the auto-covariance function $C(\mathbf{h})$. The *nugget effect* is a discontinuity in the semi-variogram for $h = 0$, which is due to both measurement uncertainties and micro variability that cannot be revealed at the scale of sampling. The *range of influence* is the distance where co-variation between samples ceases to exist; measurements taken further apart are uncorrelated.

Two other models often used are the exponential model (see Figure 4)

$$\gamma^*(h) = \begin{cases} 0 & h = 0 \\ C_0 + C_1 [1 - \exp(-\frac{3h}{R})] & h > 0 \end{cases}$$

and the Gaussian model (see Figure 5)

$$\gamma^*(h) = \begin{cases} 0 & h = 0 \\ C_0 + C_1 [1 - \exp(-\frac{3h^2}{R^2})] & h > 0. \end{cases}$$

These latter two models never reach but approach the *sill* asymptotically. Due to its horizontal tangent for $h \rightarrow 0$ the Gaussian model is good for describing very continuous phenomena.

Also other semi-variogram models such as linear and power functions are sometimes applied. To allow for so-called *nested structures* where the semi-variogram has different structures depending on the magnitude and possibly the direction of the displacement vector between observations, combinations of models may be useful.

The model parameters may be estimated b.m.o. iterative, non-linear least squares methods. These minimise the squared differences between the experimental semi-variogram and the model considered as a function of the vector of parameters θ , here $\theta = [C_0 \ C_1 \ R]^T$

$$\min_{\theta} \|\hat{\gamma}(\mathbf{h}) - \gamma^*(\theta, \mathbf{h})\|^2.$$




Fig. 1. Simple example with three observations.

For examples on an experimental semi-variogram and different models, see Figures 4 and 5.

Note, that $C(0)$ is the auto-covariance function for displacement vector $\mathbf{h} = \mathbf{0}$, and that C_0 is a parameter in the semi-variogram model.

C. Examples

To illustrate the calculations Figure 1 shows a very simple example with three observations, $z_1 = 1, z_2 = 3$ og $z_3 = 2$ with (1-D) coordinates $-2, -1$ og 3 . The semi-variogram $\hat{\gamma}$ with $\Delta h = 1.5$ is calculated like this (lags are distance groups defined by $h \pm \Delta h$)

lag	h	N	$\hat{\gamma}$
0	$0 < h \leq 3$	1	$1/2(1-3)^2 = 2$
1	$3 < h \leq 6$	2	$1/4((1-2)^2 + (3-2)^2) = 1/2$

As another more realistic example Figure 2 shows a map with sample sites. Each circle is centered on a sample point and its radius is proportional to the quantity measured, in this case the height above the ground water in a $10 \text{ km} \times 10 \text{ km}$ area near Slagelse, Denmark. Figure 3 shows a histogram for these data.

Figure 4 shows all possible pairwise squared differences as a function of distance between observations for the height data (assuming isotropy). Also an exponential variogram model estimated directly on this point cloud is shown. The *nugget effect* is 0 m^2 , the effective *range* is $3,840 \text{ m}$ and the *sill* is 840 m^2 (corresponding to 420 m^2 for the semi-variogram model).

Figure 5 shows the corresponding experimental semi-variogram. Δh is here 100 m and again we assume isotropy. Traditionally the first lag interval is half the size of the remaining lags, here 50 m . The experimental semi-variogram indicates




Fig. 2. Sample sites, each circle is centered on a sample point, radius is proportional to the quantity measured, in this case the height above the ground water in a $10 \text{ km} \times 10 \text{ km}$ area near Slagelse, Denmark.




Fig. 3. Simple statistics and histogram for height data near Slagelse, Denmark.




Fig. 4. All possible pairwise squared differences as a function of the magnitude of the displacement vector; exponential variogram model shown.




Fig. 5. Experimental semi-variogram as a function of the magnitude of the displacement vector; Gaussian semivariogram model shown.

that a Gaussian model may perform better than the exponential model in this case. Therefore a Gaussian model based on the experimental semi-variogram is shown also. The *nugget effect* is 18 m², the *range* is 1,890 m and the *sill* is 364 m².

III. SPATIAL INTERPOLATION

This section deals with deterministic types of interpolation such as inverse distance weighting and radial basis functions, and statistical types known under the joint name of kriging. Specifically equations for simple and ordinary kriging are deduced.

Other deterministic interpolation methods use (Delaunay) triangulation, Voronoi tessellation, regression analysis for determination of trend surfaces, minimum curvature etc., [15], [3].

A. Inverse Distance Weighting

Possibly the simplest conceivable way of carrying out interpolation consists of assigning the value of the nearest neighbour to a point where the value is unknown. An potential improvement consists of assigning higher weights to observations closer to the points to which we interpolate. An obvious way of doing this is to assign weights that are proportional to the inverse distance from the desired point to all N points entering into the interpolation. For the i th point we get the weight

$$w_i = \frac{1/d_i}{\sum_{j=1}^N 1/d_j},$$

where d_j is the distance from point j to the point to which we interpolate. This is readily extended to weighting with different powers, $p > 0$, of the inverse distance

$$w_i = \frac{1/d_i^p}{\sum_{j=1}^N 1/d_j^p}.$$

A.1 Examples

We now wish to interpolate to Z_0 at position $r = 0$ in Figure 1 b.m.o. inverse distance weighting. d_i is the distance from point Z_i to Z_0 . We readily calculate the following weights

r	d_i	$1/d_i$	$(1/d_i)/\sum(1/d_i)$
-2	2	1/2	3/11 (= 0.2727)
-1	1	1	6/11 (= 0.5455)
3	3	1/3	2/11 (= 0.1818)

For different powers of d_i we get the weights

r	d_i	$p = 0.1$	$p = 2.0$	$p = 10.0$
-2	2	0.3298	0.1837	0.0010
-1	1	0.3535	0.7347	0.9990
3	3	0.3167	0.0816	0.0000

We see that for low values of p the weights approach $1/N$ for all points used. For high values of p we get near a weight of one for the nearest neighbour.

B. Radial Basis Function Interpolation

Consider a linear estimate $\hat{z}_0 = \hat{z}(r_0)$ at location r_0 based on N measurements $\mathbf{z} = [z(r_1), \dots, z(r_N)]^T = [z_1, \dots, z_N]^T$ at locations $[r_1, \dots, r_N]^T$. Assume that each observation influences its surroundings in the same way in all directions and that the influence is expressed by some function ϕ (the radial basis function, RBF) which depends on the distance $h = \|r_0 - r_i\|$

between locations \mathbf{r}_0 and \mathbf{r}_i only, $\phi = \phi(\|\mathbf{r}_0 - \mathbf{r}_i\|)$. We shall look into choices of ϕ shortly. Define the interpolated value

$$\hat{z}_0 = \sum_{i=1}^N w_i \phi(\|\mathbf{r}_0 - \mathbf{r}_i\|),$$

where w_i is the weight associated with location i . Let us determine the w_i so that the interpolation becomes exact at the known locations \mathbf{r}_j , i.e.,

$$z_j = \sum_{i=1}^N w_i \phi(\|\mathbf{r}_j - \mathbf{r}_i\|), \quad j = 1, \dots, N.$$

This makes up N equations with N unknowns, the w_i , which can be written in matrix form

$$\begin{bmatrix} \phi(\|\mathbf{r}_1 - \mathbf{r}_1\|) & \cdots & \phi(\|\mathbf{r}_1 - \mathbf{r}_N\|) \\ \vdots & \ddots & \vdots \\ \phi(\|\mathbf{r}_N - \mathbf{r}_1\|) & \cdots & \phi(\|\mathbf{r}_N - \mathbf{r}_N\|) \end{bmatrix} \begin{bmatrix} w_1 \\ \vdots \\ w_N \end{bmatrix} = \begin{bmatrix} z_1 \\ \vdots \\ z_N \end{bmatrix}$$

or

$$\Phi \mathbf{w} = \mathbf{z}.$$

Often a polynomial $P(\mathbf{r}_j) = c_0 + c_1 x_j + c_2 y_j + \cdots = [1 \ x_j \ y_j \ \dots]^T \mathbf{c}$ where $\mathbf{c} = [c_0 \ c_1 \ c_2 \ \dots]^T$ is the vector of coefficients for the polynomial is added to the interpolation at location (in two dimensions) $\mathbf{r}_j = [x_j \ y_j]^T$

$$z_j = P(\mathbf{r}_j) + \sum_{i=1}^N w_i \phi(\|\mathbf{r}_j - \mathbf{r}_i\|),$$

so that $\mathbf{P}^T \mathbf{w} = \mathbf{0}$. Here

$$\mathbf{P} = \begin{bmatrix} 1 & x_1 & y_1 & x_1^2 & y_1^2 & x_1 y_1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_N & y_N & x_N^2 & y_N^2 & x_N y_N & \cdots \end{bmatrix}$$

defines the polynomial applied. Often a constant corresponding to the column of ones or a linear polynomial is used.

Solving the system of equations $\Phi \mathbf{w} = \mathbf{z}$ under the constraint $\mathbf{P}^T \mathbf{w} = \mathbf{0}$ we get (see above on ordinary kriging and the Lagrange multiplier technique)

$$\begin{bmatrix} \Phi & \mathbf{P} \\ \mathbf{P}^T & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{w} \\ \mathbf{c} \end{bmatrix} = \begin{bmatrix} \mathbf{z} \\ \mathbf{0} \end{bmatrix}.$$

The order of the polynomial may depend on the RBF chosen.

References on RBF are [16], [17], [18].

B.1 Normalized RBF Interpolation

RBF interpolation comes in a normalized version also

$$\hat{z}_0 = \frac{\sum_{i=1}^N w_i \phi(\|\mathbf{r}_0 - \mathbf{r}_i\|)}{\sum_{i=1}^N \phi(\|\mathbf{r}_0 - \mathbf{r}_i\|)}$$

which leads to

$$z_j \sum_{i=1}^N \phi(\|\mathbf{r}_j - \mathbf{r}_i\|) = \sum_{i=1}^N w_i \phi(\|\mathbf{r}_j - \mathbf{r}_i\|), \quad j = 1, \dots, N.$$

for the weights. In matrix form this leads to the same matrix equation as above with the elements in the vector on the right hand side changed from z_j to $z_j \sum_{i=1}^N \phi(\|\mathbf{r}_j - \mathbf{r}_i\|)$.

B.2 Choice of RBF

Often one of the following choices is made for ϕ

- multiquadric: $\phi(h) = (h^2 + h_0^2)^{1/2}$,
- inverse multiquadric: $\phi(h) = (h^2 + h_0^2)^{-1/2}$,
- thin-plate spline: $\phi(h) = h^2 \log(h/h_0)$ (which tends to 0 for h tending to 0), or
- Gaussian: $\phi(h) = \exp(-\frac{1}{2}(h/h_0)^2)$,

where h_0 is a scale parameter to be chosen. Generally, h_0 should be chosen larger than a typical distance between samples and smaller than the size of the study area. The multiquadric is said to be less sensitive to the choice of h_0 . Especially the Gaussian is sensitive to this choice.

With a Gaussian RBF you don't need the polynomials mentioned above, with the thin-plate spline RBF a linear polynomial may be needed.

B.3 Shepard Interpolation

A special case for the normalized RBF interpolation consists of setting the matrix on the left hand side to a constant times the identity matrix. This corresponds to applying a ϕ that tends to infinity for h tending to 0, and is finite for $h > 0$. This leads to setting the weights equal to the measurements $w_i = z_i$, i.e., we needn't solve the system of equations for \mathbf{w} . In this case $\phi(h) = h^{-p}$, $1 < p \leq 3$ with appropriate handling for $h = 0$ is often used.

C. Kriging

Kriging (after the South African mining engineer and professor Danie Krige) is a name for a family of methods for minimum error variance estimation. Consider a linear (or rather affine) estimate $\hat{z}_0 = \hat{z}(\mathbf{r}_0)$ at location \mathbf{r}_0 based on N measurements $\mathbf{z} = [z(\mathbf{r}_1), \dots, z(\mathbf{r}_N)]^T = [z_1, \dots, z_N]^T$

$$\hat{z}_0 = w_0 + \sum_{i=1}^N w_i z_i = w_0 + \mathbf{w}^T \mathbf{z},$$

where w_i are the weights applied to z_i and w_0 is a constant.

We consider z_i as realisations of stochastic variables Z_i , $\mathbf{Z} = [Z(\mathbf{r}_1), \dots, Z(\mathbf{r}_N)]^T = [Z_1, \dots, Z_N]^T$. We think of $Z(\mathbf{r})$ as consisting of a mean value and a residual $Z(\mathbf{r}) = \mu(\mathbf{r}) + \epsilon(\mathbf{r})$ with mean value zero and constant variance σ^2 , $E\{\epsilon\} = 0$ and $\text{Var}\{\epsilon\} = \sigma^2$. For the linear estimator we get

$$\hat{Z}_0 = w_0 + \mathbf{w}^T \mathbf{Z}. \quad (2)$$

The estimation error $z_0 - \hat{z}_0$ is unknown. But for the expectation value of the estimation error we get

$$\begin{aligned} E\{Z_0 - \hat{Z}_0\} &= E\{Z_0 - w_0 - \mathbf{w}^T \mathbf{Z}\} \\ &= \mu_0 - w_0 - \mathbf{w}^T \boldsymbol{\mu}, \end{aligned} \quad (3)$$

where $\mu_0 = \mu(\mathbf{r}_0)$ is the expectation value of Z_0 and $\boldsymbol{\mu}$ is a vector of expectation values for \mathbf{Z}

$$\boldsymbol{\mu} = \begin{bmatrix} \mu(\mathbf{r}_1) \\ \vdots \\ \mu(\mathbf{r}_N) \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \vdots \\ \mu_N \end{bmatrix}.$$

We want our estimator to be unbiased or central, i.e., we demand $E\{Z_0 - \hat{Z}_0\} = 0$ or

$$\mu_0 - w_0 - \mathbf{w}^T \boldsymbol{\mu} = 0. \quad (4)$$

The variance of the estimation error is

$$\begin{aligned}\sigma_E^2 &= \text{Var}\{Z_0 - \hat{Z}_0\} \\ &= \text{Var}\{Z_0\} + \text{Var}\{w_0 + \mathbf{w}^T \mathbf{Z}\} \\ &\quad - 2 \text{Cov}\{Z_0, w_0 + \mathbf{w}^T \mathbf{Z}\} \\ &= \sigma^2 + \mathbf{w}^T (\mathbf{Cw} - 2 \text{Cov}\{Z_0, \mathbf{Z}\}),\end{aligned}$$

where \mathbf{C} is the dispersion or variance-covariance matrix of the stochastic variables, \mathbf{Z} , entering into the estimation.

What is said in Section III-C so far is valid for all linear estimators. The idea in kriging is now to find the linear estimator which minimises the estimation variance.

C.1 Simple Kriging

In simple kriging (SK) we assume that $\mu(\mathbf{r})$ is known. From Equations 2 and 4 we get

$$\hat{Z}_0 - \mu_0 = \mathbf{w}^T (\mathbf{Z} - \boldsymbol{\mu}).$$

The weights w_i are found by minimising the estimation variance σ_E^2 . This is done by setting the partial derivatives to zero

$$\frac{\partial \sigma_E^2}{\partial \mathbf{w}} = 2 \mathbf{Cw} - 2 \text{Cov}\{Z_0, \mathbf{Z}\} = \mathbf{0},$$

which results in the SK system

$$\mathbf{Cw} = \text{Cov}\{Z_0, \mathbf{Z}\}$$

or

$$\begin{bmatrix} C_{11} & \cdots & C_{1N} \\ \vdots & \ddots & \vdots \\ C_{N1} & \cdots & C_{NN} \end{bmatrix} \begin{bmatrix} w_1 \\ \vdots \\ w_N \end{bmatrix} = \begin{bmatrix} C_{01} \\ \vdots \\ C_{0N} \end{bmatrix},$$

where C_{ij} , $i, j = 1, \dots, N$ is the covariance between points i and j among the N points, which enter into the estimation of point 0. C_{0j} , $j = 1, \dots, N$ is the covariance between point j and point 0, the point to which we interpolate. We get these covariances from the semi-variogram model (remembering Equation 1, $\gamma(\mathbf{h}) = C(\mathbf{0}) - C(\mathbf{h})$) as the sill minus the value of the semi-variogram model for the relevant distance (and possibly direction) between observations. (Alternatively, the kriging system may be formulated b.m.o. the semi-variogram; to avoid zeros on the diagonal of \mathbf{C} we prefer the covariance formulation for numerical reasons.) Here C_{ij} must not be confused with the semi-variogram parameters C_0 and C_1 .

The minimised squared estimation error termed the simple kriging variance is

$$\begin{aligned}\sigma_{\text{SK}}^2 &= \sigma^2 + \mathbf{w}^T (\mathbf{Cw} - 2 \text{Cov}\{Z_0, \mathbf{Z}\}) \\ &= \sigma^2 - \mathbf{w}^T \text{Cov}\{Z_0, \mathbf{Z}\}.\end{aligned}$$

In SK the mean value $\mu(\mathbf{r})$ is known. In practice it is often assumed constant for the entire domain (or study area), or we must estimate it before the interpolation (or we must construct an interpolation algorithm which does not require knowledge of the mean field, see the next section).

C.2 Ordinary Kriging

In ordinary kriging (OK) we assume that the mean $\mu(\mathbf{r})$ is constant and equal to μ_0 for Z_0 and the N points that enter into the estimation of Z_0 . From Equations 3 and 4 we get

$$\mathbb{E}\{Z_0 - \hat{Z}_0\} = \mu_0(1 - \mathbf{w}^T \mathbf{1}) - w_0 = 0$$

for any μ_0 . $\mathbf{1}$ is a vector of ones. This is possible only if $w_0 = 0$ and $\mathbf{w}^T \mathbf{1} = 1$.

The weights w_i are found by minimising σ_E^2 under the constraint $\mathbf{w}^T \mathbf{1} = 1$. A standard technique for minimisation under a constraint is introducing a function F with a so-called Lagrange multiplier (here -2λ) which we multiply by the constraint set to zero and then minimising

$$F = \sigma_E^2 + 2\lambda(\mathbf{w}^T \mathbf{1} - 1)$$

without constraints. Again the partial derivatives are set to zero

$$\begin{aligned}\frac{\partial F}{\partial \mathbf{w}} &= 2 \mathbf{Cw} - 2 \text{Cov}\{Z_0, \mathbf{Z}\} + 2\lambda \mathbf{1} = \mathbf{0} \\ \frac{\partial F}{\partial \lambda} &= 2(\mathbf{w}^T \mathbf{1} - 1) = 0,\end{aligned}$$

which results in the OK system

$$\begin{aligned}\mathbf{Cw} + \lambda \mathbf{1} &= \text{Cov}\{Z_0, \mathbf{Z}\} \\ \mathbf{1}^T \mathbf{w} &= 1\end{aligned}$$

or

$$\begin{bmatrix} C_{11} & \cdots & C_{1N} & 1 \\ \vdots & \ddots & \vdots & \vdots \\ C_{N1} & \cdots & C_{NN} & 1 \\ 1 & \cdots & 1 & 0 \end{bmatrix} \begin{bmatrix} w_1 \\ \vdots \\ w_N \\ \lambda \end{bmatrix} = \begin{bmatrix} C_{01} \\ \vdots \\ C_{0N} \\ 1 \end{bmatrix}.$$

The values requested for C_{ij} are found as described in the previous section on SK.

The minimised squared estimation error termed the ordinary kriging variance is

$$\begin{aligned}\sigma_{\text{OK}}^2 &= \sigma^2 + \mathbf{w}^T (\mathbf{Cw} - 2 \text{Cov}\{Z_0, \mathbf{Z}\}) \\ &= \sigma^2 - \mathbf{w}^T \text{Cov}\{Z_0, \mathbf{Z}\} - \lambda.\end{aligned}$$

OK implies an implicit re-estimation of μ_0 for each new constellation of points. This is an attractive property making OK well suited for interpolation in situations where the mean is not constant (i.e., in the absence of first order stationarity).

C.3 Examples

Let us consider the data in Figure 1 again. We now wish to interpolate to the position $r = 0$ b.m.o. ordinary kriging. To carry out the calculations we use a stipulated semi-variogram based on the spherical model with $C_0 = 0$, $C_1 = 1$ and $R = 6$. Remembering Equation 1, $C(h) = C(\mathbf{0}) - \gamma(h)$, this gives the auto-covariance function (in this case where $C_0 + C_1 = 1$ this is the same as the auto-correlation function)

h	$\hat{\gamma}(h)$	$C(h)$
0	0.0000	1.0000
1	0.2477	0.7523
2	0.4815	0.5185
3	0.6875	0.3125
4	0.8519	0.1481
5	0.9606	0.0394
6	1.0000	0.0000

Therefore the OK system looks like this

$$\begin{bmatrix} 1.0000 & 0.7523 & 0.0394 & 1 \\ 0.7523 & 1.0000 & 0.1481 & 1 \\ 0.0394 & 0.1481 & 1.0000 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ \lambda \end{bmatrix} = \begin{bmatrix} 0.5185 \\ 0.7523 \\ 0.3125 \\ 1 \end{bmatrix}$$




Fig. 6. Kriged map of heights above the ground water (unit is m).




Fig. 7. Kriging variance corresponding to Figure 6 (unit is m^2).

where the values for C_{ij} come from the $C(h)$ table. The solution is $w_1 = -0.0407$, $w_2 = 0.7955$, $w_3 = 0.2452$ and $\lambda = -0.0489$, which gives a kriging variance of 0.3949. We see that even though Z_1 is closer to Z_0 than Z_3 , the weight on Z_1 is much smaller than the weight on Z_3 . This is an attractive characteristic of kriging, which allows for possible clustering of the sampling locations. We say that Z_2 *screens* for Z_1 . This *screening* effect becomes weaker for higher nugget effects and it disappears for pure nugget effect (i.e., $C_1 = 0$ for the models shown here), where all weights become equal.

In a more realistic example Figure 6 shows a kriged (OK) map over heights above the ground water for the area near Slagelse. The interpolation is based on the isotropic Gaussian model for the experimental semi-variogram in Figure 5 (nugget effect 18 m^2 , range 1.890 m and sill 364 m^2). We have kriged to 100 by 100 points in a 100 m by 100 m grid using a moving window to include the points from which to interpolate. The search radius of the moving window was 2,000 m and the estimation for each point was based on minimum 4 and maximum 20 points. We see that the interpolated map shows a good correspondance with the map of sample sites in Figure 2. Figure 7 shows the corresponding OK variances. We see that the kriging variance is large where the distances to the nearest samples are large.

C.4 Other Types of Kriging

If we wish to estimate average (also known as regularised) values over an area or a volume rather than point values, we may use *block kriging* which can be combined with several other forms of kriging.

If more variables are studied simultaneously the above methods for description of spatial correlation may be extended to handle the spatial covariation between all pairs of variables in the form of cross semi-variograms or cross-covariance functions. Also more variables may be interpolated simultaneously using *cokriging*. *Cokriging* is most useful when one variable is sampled on fewer locations than other correlated variables. *Universal kriging* is a method for the case where the mean value is described by linear combinations of known functions ideally determined by the physics of the problem at hand. Also methods for non-linear kriging exist such as *lognormal kriging*, *multi-Gaussian kriging*, *rank kriging*, *indicator kriging* and *disjunctive kriging*. References here are [1], [3], [9], [14].

C.5 Conclusions, kriging

The above sections and examples demonstrate the following properties of kriging:

- Kriging is a type of interpolation that gives us both an estimate based on the spatial structure of the variable in question as expressed by the auto-covariance function (or the semi-variogram) as well as an estimation variance which is minimised.
- The kriging estimator is the best linear unbiased estimator (BLUE) in the sense that it minimises the estimation variance. Also it is exact, i.e., if we interpolate to a point which coincides with an existing sample point, kriging gives the same value as the one measured and the kriging variance is zero.
- The kriging system and the kriging variance depend on the auto-covariance function (or the semi-variogram) and the spatial layout of the sample locations only and not on the actual data values. If an auto-covariance function (or a semi-variogram) is known (or assumed or imposed) this has important potential for minimising the estimation variance in experimental design (i.e., in the planning phase of the spatial lay-out of the sampling scheme).
- The solution of the kriging system implies a statistical distance weighting of the data values which enter into the interpolation. Also for OK, the weights are scaled so they add to one. Furthermore, possible redundancy in the form of clustering of the sample locations is accounted for; the above mentioned screening effect is due to this de-clustering.
- Because of the implicit re-estimation of the mean value for each new point constellation, OK is well suited for situations where the mean is not constant over the study region, i.e., where we don't have first order stationarity.

Further, the kriging system has a unique solution if and only if the covariance matrix \mathbf{C} (Section III-C) is positive definite; this also guarantees a non-negative kriging variance.

The strength of kriging may be attributed to a combination of the above characteristics.

If we choose to formulate the kriging system in terms of the auto-covariance function which is done in this note, we must assume (or impose) second order stationarity, i.e., the same auto-covariance function over the entire study area. The system may also be formulated in terms of the semi-variogram and in this case we must assume the intrinsic hypothesis, i.e., the same semi-variogram over the entire study area.

These assumptions may seem to be a drawback of kriging but if deterministic methods are applied, we implicitly make similar

assumptions. It can hardly be considered as a drawback of geostatistical methods that we are forced to consider whether such assumptions are appropriate.

IV. FINAL REMARKS

A sampling strategy may be based on the dependence of the kriging variance on the distance to the nearest samples. If the auto-covariance function (or the semi-variogram) and the sample locations are known, we can determine the kriging weights and the kriging variances before the actual sampling takes place. If the variances become too large in some regions of our study area we may modify the sample locations to obtain smaller variances. Also, to obtain a good estimate of the nugget effect which is an important parameter for the outcome of the kriging process, it may be an advantage to position some samples close to each other.

In multivariate studies where the joint behaviour of several variables is investigated, rather than interpolating the original variables we may interpolate combinations of them. For instance we may interpolate principal components or factors resulting from a factor analysis or a spatial factor analysis, [19], [20], [11], [21], [22]. General references to multivariate statistics are for example [23], [24]. [25] is written especially for geographers, [15] for geologists.

Also temporal aspects in connection with the application of data that vary in both space and time may be important. Spatio-temporal semi-variograms and spatio-temporal kriging are dealt with in for example [26], [27]. A GIS for handling of temporal data is described in [28].

REFERENCES

- [1] A. G. Journel and Ch. J. Huijbregts, *Mining Geostatistics*, Academic Press, London, 1978, 600 pp.
- [2] Isobel Clark, *Practical Geostatistics*, Elsevier Applied Science, London, 1979, <http://www.stokos.demon.co.uk/>.
- [3] Edward H. Isaaks and R. Mohan Srivastava, *An Introduction to Applied Geostatistics*, Oxford University Press, New York, 1989, 561 pp.
- [4] K. Conradsen, A. A. Nielsen, and K. Windfeld, "Analysis of geochemical data sampled on a regional scale," in *Statistics in the Environmental and Earth Sciences*, A. Walden and P. Guttorp, Eds., pp. 283–300. Griffin, 1992.
- [5] Noel A. C. Cressie, *Statistics for Spatial Data*, revised edition, Wiley, New York, 1993.
- [6] P. A. Burrough and R. A. MacDonnell, *Principles of Geographical Information Systems*, Oxford University Press, 1998.
- [7] M. F. Goodchild, B. O. Parks, and L. T. Steyaert, *Environmental Modeling with GIS*, Oxford University Press, 1993.
- [8] H. Wackernagel, *Multivariate Geostatistics*, third edition, Springer, 2003.
- [9] C. V. Deutsch and A. G. Journel, *GSLIB: Geostatistical Software Library and User's Guide*, second edition, Oxford University Press, 1998, Internet <http://www.gslib.com/>.
- [10] Y. Pannatier, *VARIOWIN: Software for Spatial Data Analysis in 2D*, Springer, 1996, Internet <http://www-sst.unil.ch/research/variowin/>.
- [11] Allan A. Nielsen, *Analysis of Regularly and Irregularly Sampled Spatial, Multivariate, and Multi-temporal Data*, Ph.D. thesis, Informatics and Mathematical Modelling, Technical University of Denmark, Lyngby, 1994, Internet <http://www.imm.dtu.dk/pubdb/p.php?296>.
- [12] Karsten Hartelius, *Analysis of Irregularly Distributed Points*, Ph.D. thesis, Informatics and Mathematical Modelling, Technical University of Denmark, Lyngby, 1996. Internet <http://www.imm.dtu.dk/pubdb/p.php?1204>.
- [13] Allan A. Nielsen, "2D semivariograms," in *Proceedings of the Fourth South African Workshop on Pattern Recognition*, Paul Cilliers, Ed., Simon's Town, South Africa, 25–26 November 1993, pp. 25–35.
- [14] Allan A. Nielsen, "Kriging," lecture note, Technical University of Denmark, 2004. Internet <http://www.imm.dtu.dk/pubdb/p.php?3479>.
- [15] J. C. Davis, *Statistics and Data Analysis in Geology*, second edition, John Wiley & Sons, 1986.
- [16] M. J. D. Powell, "The theory of radial basis function approximation," in *Advances in Numerical Analysis II: Wavelets, Subdivision Algorithms and Radial Functions*, W. A. Light, Ed., Oxford University Press, 1992.
- [17] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes: The Art of Scientific Computing*, third edition, Cambridge University Press, 2007.
- [18] F. Anton, J. A. Bærentzen, J. Gravesen, and H. Aanæs, *Computational Geometry Processing*, lecture note, Technical University of Denmark, 2008.
- [19] E. C. Grunsky and F. P. Agterberg, "Spatial and multivariate analysis of geochemical data from metavolcanic rocks in the Ben Nevis area, Ontario," *Mathematical Geology*, vol. 20, no. 7, pp. 825–861, 1988.
- [20] E. C. Grunsky and F. P. Agterberg, "SPFAC: a Fortran program for spatial factor analysis of multivariate data," *Computers & Geosciences*, vol. 17, no. 1, pp. 133–160, 1991.
- [21] Allan A. Nielsen, Knut Conradsen, John L. Pedersen, and Agnæte Steenfelt, "Spatial factor analysis of stream sediment geochemistry data from South Greenland," in *Proceedings of the Third Annual Conference of the International Association for Mathematical Geology*, Vera Pawlowsky-Glahn, Ed., Barcelona, Spain, September 1997, pp. 955–960.
- [22] Allan A. Nielsen, Knut Conradsen, John L. Pedersen, and Agnæte Steenfelt, "Maximum autocorrelation factorial kriging," in *Proceedings of the 6th International Geostatistics Congress (Geostats 2000)*, W. J. Kleingeld and D. G. Krige, Eds., Cape Town, South Africa, April 2000. Internet <http://www.imm.dtu.dk/pubdb/p.php?3639>.
- [23] Knut Conradsen, *Introduktion til Statistik*, Informatics and Mathematical Modelling, Technical University of Denmark, 1984.
- [24] T. W. Anderson, *An Introduction to Multivariate Statistical Analysis*, second edition, John Wiley, New York, 1984.
- [25] D. A. Griffith and C. G. Amrhein, *Multivariate Statistical Analysis for Geographers*, Prentice Hall, 1997.
- [26] Annette Kjær Ersbøll, "A comparison of two spatio-temporal semivariograms with use in agriculture," in *Lecture Notes in Statistics*, T. G. Gregoire et al., Ed., vol. 122, pp. 299–308. Springer-Verlag, 1997.
- [27] Annette Kjær Ersbøll and Bjarne Kjær Ersbøll, "On spatio-temporal kriging," in *Proceedings of the Third Annual Conference of the International Association for Mathematical Geology (IAMG'97)*, Vera Pawlowsky-Glahn, Ed., Barcelona, Spain, September 1997, pp. 617–622.
- [28] Thomas Knudsen, *Busstop - A Spatio-Temporal Information System*, Ph.D. thesis, Niels Bohr Institute, Department of Geophysics, University of Copenhagen, Denmark, 1997.

INDEX TERMS

auto-correlation function
auto-covariance function
BLUE
data analysis
distance weighting
estimation variance
geostatistics
GIS
GSLIB
intrinsic hypothesis
covariance function
kriging
kriging variance
Lagrange multiplier
least squares method
multivariate statistics
nested structures
normalized radial basis functions
nugget effect
ordinary kriging
sampling strategy
radial basis functions
range of influence
regionalised variable
screening effect
semi-variogram
semi-variogram model
sill
simple kriging
spatial interpolation
spatial correlation
stationarity
statistics
stochastic function
stochastic process
stochastic variable
variogram
Variowin