


Seventeen years of global SSH anomalies analyzed by a maximum information based extension to EOF analysis



Jacob S. Vestergaard, Allan A. Nielsen, Ole B. Andersen
DTU Space, Technical University of Denmark

Introduction

Empirical Orthogonal Function (EOF) is traditionally used for analyzing sea level data as it provides a compact description of the spatial and temporal variability in terms of orthogonal functions. The motivation for extending the Empirical Orthogonal Function (EOF) analysis to an information maximization based approach, is to find descriptive subspaces where a large amount of information is present.

17 years of global Sea Surface Height (SSH) anomalies is analyzed using the Infomax algorithm as the application of this method can identify latent oceanographic features in the data set, with no other constraints than independence.


Expected to be prominent in the analyses is the El Niño Southern Oscillation (ENSO), which is a combined oceanographic and atmospheric phenomenon. The occurrence of El Niño is periodic with an inexact cycle time of 2-7 years and has enormous socio-economic consequences all over the planet. The ENSO consists of different

Data

The data set are 17 years of global Sea Surface Height (SSH) anomalies from TOPEX/Poseidon, Jason-1 and Jason-2 missions.

Period: January 1993 → December 2009 (204 months).

Arranged in a grid of size 67×120 with resolution of $2^\circ \times 3^\circ$.



The $P = 204$ months of SSH values are represented in the data matrix X with the months column wise and $N = 8040$ observations row wise.

$$X = \begin{bmatrix} x_{1,1} & \dots & x_{1,P} \\ \vdots & \ddots & \vdots \\ x_{N,1} & \dots & x_{N,P} \end{bmatrix} = [x_1 \ x_2 \ \dots \ x_P] \quad (1)$$

Prior to these analyses, the data have been preprocessed to extract seasonal signals, tide, Earth tide and impacts from high/low pressure weather systems.




Figure 1: Mean values and first order fit. Mean increase per year approximately 2.8729 mm

EOF analysis - a summary

Empirical Orthogonal Function (EOF) analysis is a variant of Principal Component Analysis (PCA), where the assumption of zero spatial mean is replaced by an assumption of zero temporal mean.

To find the principal directions, the empirical dispersion matrix is calculated as

$$\hat{\Sigma} = \frac{1}{N-1} X^T X \quad (2)$$

which is recognized as T-mode analysis [3]. The principal components (EOFs) are calculated by finding the the eigenvectors of $\hat{\Sigma}$ that satisfy

$$\hat{\Sigma} u_j = \lambda_j u_j \quad (3)$$

followed by a projection of the original data onto these

$$EOF_j = X u_j \quad \forall j \in [1, P] \quad (4)$$

The EOFs are usually sorted decreasing according to contained variance.

Results The five most significant EOFs and loadings are displayed in Figure 2. All have a large peak with relation to El Niño of 1997-1998. Not identical - separated by approx. one month.




Figure 2: First five EOFs and loadings. All dominated by El Niño 1997-1998 in different phases.

Maximizing Information (Infomax)

The Information Maximization (Infomax [1]) method is part of the ICA family of methods, which seek to uncover a number of latent sources. The sources s_i have been put through a mixing matrix G and intercepted as signals in X

$$X^T = GS \quad (5)$$

where $S = [s_1 \ s_2 \ \dots \ s_P]^T$.

Sources are sought to contain maximum information, which corresponds to nongaussianity [2]. No constraint of being mutual orthogonal.

The Infomax method is a maximum likelihood based approach to finding the directions in space that maximize nongaussianity, using e.g. steepest ascent. The update rule for adjusting the weights w is

$$\Delta W \propto W^T + z z^T \quad (6)$$

where $z = [\phi(w_1^T x) \ \phi(w_2^T x) \ \dots \ \phi(w_P^T x)]$, with $\phi(w^T x) = -\tanh(w^T x) \Rightarrow$ implicit assumption that the sources are leptokurtic (or supergaussian).




Figure 3: Assumed distribution and Gaussian




Figure 4: First five ICs and corresponding loadings

Results The first five independent components (ICs) can be identified to contain the following information, from the temporal and spatial patterns in Figure 4:

1. A large pool of water hits South America from the west. Large peak at El Niño of 1997-1998, and peaks at other known El Niño years.
2. Captures periods following El Niños - part of the build-up phase to the next.
3. The southern Pacific is left colder than usual (La Niña).
4. Strong linear trend in the temporal pattern. The spatial representation highlights - among others - an area south of Greenland.
5. Captures a connection between El Niños and the SSH anomalies in the Indian Ocean.

Further analysis

IC1's El Niño related peaks in the temporal loadings of have been marked in Figure 5.





Figure 5: Period estimation from IC1

A Fourier analysis has been used to determine the periodicity of El Niño in the time series. Due to few occurrences no robust estimates are possible.

Highlighted points south of Greenland in IC4 and mean SSH anomalies for these in Figure 6.



A stronger trend than the global average through majority of period.




Figure 6: SSH anomalies in area highlighted in IC4

Trend started turning in 2006-2007?

Conclusions

Main results

1. Detection of a sea level rise south of Greenland through the majority of the time period. The information extracted from this data set, however, suggests that the trend started turning in 2006-2007, similarly to 1999-2001.
2. Purer separation of the ENSO signal in the first component. Pure separation of signals of interest simplifies further analysis and hypothesis generation or confirmation. The periodicity of El Niño through the last 17 years have been estimated, without much success, though, due to short time period.

Infomax pros and cons

- + No demand for mutual orthogonality.
- + Flexibility by allowing for different nonlinear mappings and thereby different assumptions about the source priors.
- Assumption of equal number of sources and intercepted signals.
- Iterative optimization.

Overall, when analyzing these 17 years of SSH anomalies, Infomax has shown superior to EOF analysis in extracting information of interest.

References

- [1] A. Bell and T.J. Sejnowski. An information-maximization approach to blind separation and blind deconvolution. *Neural Computation*, 7:1129-1159, 1995.
- [2] Aapo Hyvärinen, Juha Karhunen, and Erkki Oja. *Independent Component Analysis*. Wiley-Interscience, May 2001.
- [3] Michael B. Richman. Rotation of principal components. *International Journal of Climatology*, 6(3):293-335, 1986.